# QSAR model for CYP450 3A4 inhibitor (v1.0)



### **ProtoADME**

ProtoADME is a computational (in silico) tool focused on the prediction of endpoints related with the ADME (Absorption, Distribution, Metabolism and Excretion) of chemical substances.

## **Endpoint**

Toxicokinetic: CYP450 3A4 inhibitor

The microsomal cytochrome P450 (CYP) family 4 monooxygenases are the major fatty acid omega-hydroxylases. These enzymes remove excess free fatty acids to prevent lipotoxicity, catabolize leukotrienes and prostanoids, and also produce bioactive metabolites from arachidonic acid omega-hydroxylation. In addition to endogenous substrates, recent evidence indicates that CYP4 monooxygenases can also metabolize xenobiotics, including therapeutic drugs. If a compound is a CYP inhibitor may decrease the metabolism of comedicated drugs.

## **Metrics**

#### **Training set**

QSAR predictions		
Non-inhibitor	Inhibitor	
249	5	
30	203	
	Non-inhibitor	

<b>\/</b> ^	liم	24	ion	COF
٧a	HU	ıαι	IUII	set

Experimental values	QSAR predictions		
	Non-inhibitor	Inhibitor	
Non-inhibitor	64	30	
Inhibitor	29	43	

Parameters	Training	Validation
Accuracy	0.93	0.64
Sensitivity / recall	0.87	0.60
Specificity	0.98	0.68
Precision	0.98	0.59
Negative predictive value	0.89	0.69
F-score	0.92	0.59
Matthews Correlation Coefficient	0.86	0.28
Critical Success Index	0.85	0.42
Area under the ROC	0.93	0.64

ProtoADME is part of



ProtoPRED platform allows the easy, fast and user-friendly prediction of different properties of chemical compounds, using proprietary (Q)SAR models.



+34 962 021 811



protopred@protogsar.com